

<https://helda.helsinki.fi>

---

## Diel- and temperature-driven variation of leaf dark respiration rates and metabolite levels in rice

Rashid, Fatimah Azzahra Ahmad

2020-10

---

Rashid , F A A , Scafaro , A P , Asao , S , Fenske , R , Dewar , R , Masle , J , Taylor , N L & Atkin , O K 2020 , ' Diel- and temperature-driven variation of leaf dark respiration rates and metabolite levels in rice ' , New Phytologist , vol. 228 , no. 1 , pp. 56-69 . <https://doi.org/10.1111/nph.16661>

---

<http://hdl.handle.net/10138/329958>

<https://doi.org/10.1111/nph.16661>

---

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# **Diel and temperature driven variation of leaf dark respiration rates and metabolite levels in rice**

Fatimah Azzahra Ahmad Rashid<sup>1,2</sup>, Andrew P. Scafaro<sup>1\*</sup>, Shinichi Asao<sup>1</sup>, Ricarda Fenkse<sup>3</sup>, Roderick C. Dewar<sup>4,5</sup>, Josette Masle<sup>4</sup>, Nicolas L. Taylor<sup>3#</sup>, Owen K. Atkin<sup>1#</sup>

<sup>1</sup> Australian Research Council Centre of Excellence in Plant Energy Biology, Research School of Biology, The Australian National University, Canberra, ACT, 2601, Australia

<sup>2</sup> Current address: Department of Biology, Faculty of Science and Mathematics, Sultan Idris education University, 35900 Tanjung Malim, Perak, Malaysia

<sup>3</sup> Australian Research Council Centre of Excellence in Plant Energy Biology, School of Molecular Sciences and Institute of Agriculture, Faculty of Science, The University of Western Australia, Crawley WA 6009, Australia

<sup>4</sup> Research School of Biology, The Australian National University, Canberra, ACT, 2601, Australia

<sup>5</sup> Institute for Atmospheric and Earth System Research/Physics, University of Helsinki, Helsinki, Finland

# The support of the Australian Research Council (ARC) Centre of Excellence in Plant Energy Biology (CE140100008) is acknowledged

\*Correspondence to: [andrew.scafaro@anu.edu.au](mailto:andrew.scafaro@anu.edu.au), Tel: +61 2 6125 9580, Fax: +61 2 6125 5095

**Keywords:** amino acids, diel cycle, growth temperature, leaf dark respiration, metabolites, rice, sugars, tricarboxylic acid (TCA) cycle

Total word count (6054); Introduction (1520); Materials and Methods (1173); Results (1196); Discussion (2165); Tables (1); Figures (6- all in colour), Supporting Information (4)

## Summary

- Leaf respiration in the dark ( $R_{\text{dark}}$ ) is often measured at a single time during the day, with hot-acclimation lowering  $R_{\text{dark}}$  at a common measuring temperature. However, it is unclear whether the diel cycle influences the extent of thermal acclimation of  $R_{\text{dark}}$ , or how temperature and time of day interact to influence respiratory metabolites.
- To examine these issues, we grew rice at 25/20°C (day/night), 30/25°C and 40/35°C, measuring  $R_{\text{dark}}$  and changes in metabolites at five time-points spanning a single 24-hour period.
- $R_{\text{dark}}$  differed among the treatments and with time of day. However, there was no significant interaction between time and growth temperature, indicating that the diel cycle does not alter thermal acclimation of  $R_{\text{dark}}$ . Amino acids were highly responsive to the diel cycle and growth temperature, and many were negatively correlated with carbohydrates and with organic acids of the tricarboxylic acid (TCA) cycle. Organic TCA intermediates were significantly altered by the diel cycle irrespective of growth temperature, which we attribute to light-dependent regulatory control of TCA enzyme activities.
- Collectively, our study shows that environmental disruption of the balance between respiratory substrate supply and demand is corrected for by shifts in TCA-dependent metabolites.

## Introduction

Leaf respiration in the dark ( $R_{\text{dark}}$ ) provides plants with ATP, reducing equivalents, and carbon skeletons needed to support growth and cellular maintenance (Atkin & Macherel, 2009, Gonzalez-Meler *et al.*, 2004, Lambers & Ribas-Carbo, 2005, Noguchi & Yoshida, 2008). Rates of  $R_{\text{dark}}$  increase exponentially with rising temperature in the short-term (i.e. time scales of seconds to minutes) (Amthor, 2000, Atkin *et al.*, 2005b, Reich *et al.*, 2016, Smith & Dukes, 2013). With sustained exposure to different growth temperatures (e.g. over periods of days to weeks),  $R_{\text{dark}}$  often acclimates, resulting in lower rates when measured at a common measuring temperature in warm/hot-grown plants compared to their cooler-grown counterparts (Atkin *et al.*, 2005a, Reich *et al.*, 2016, Slot & Kitajima, 2015, Smith & Dukes, 2017). Acclimation of

$R_{\text{dark}}$  is common (Slot & Kitajima, 2015), and likely contributes to global patterns in  $R_{\text{dark}}$  (Atkin *et al.*, 2015, Vanderwel *et al.*, 2015), with acclimation having the potential to dampen the effects of future climate warming on plant respiratory carbon release (Huntingford *et al.*, 2017, Reich *et al.*, 2016, Smith & Dukes, 2013). Importantly, studies assessing acclimation invariably rely on measurements made at a single time-point in a day, implicitly assuming that acclimation capacity is constant through a 24-hour diel cycle. This assumption remains untested, even though  $R_{\text{dark}}$  supports leaf metabolism which varies greatly during the day. For example, photosynthesis varies with diel fluctuations in light intensity, temperature, and evaporative demand, likely affecting carbon fluxes in cellular metabolism (Steer, 1973). Given the strong coupling between respiratory and photosynthetic metabolism (Kromer, 1995, Shameer *et al.*, 2019, Tcherkez *et al.*, 2017), diel and growth-temperature dependent variations in photosynthetic activity might be expected to influence rates of  $R_{\text{dark}}$ . Time of day is also likely to influence concentrations of metabolites involved in glycolysis, the tricarboxylic acid (TCA) cycle, the shikimate pathway and several other related metabolic networks (Gibon *et al.*, 2009, Hurry *et al.*, 2005, Urbanczyk-Wochniak *et al.*, 2005); however, whether diel variations in these metabolites is affected by growth temperature remains unclear. This interaction between the diel cycle and respiratory responses to growth temperature is important in understanding the effects of climate warming on plant carbon fluxes, both in natural and managed ecosystems. This is especially important when considering the disproportionate increase in night temperatures over the past half century (Davy *et al.* 2017). The impact of growth temperature and time of day on leaf respiratory metabolism is of particular interest in globally important crops such as rice, where rising growth temperatures (particularly during the night) are leading to reduced yields (Peng *et al.*, 2004, Welch *et al.*, 2010).

Photosynthesis generates sugar substrates for respiration, but the evidence for substrate-dependent changes in leaf  $R_{\text{dark}}$  is inconclusive. In some studies, glucose and sucrose concentrations can positively correlate with rates of  $R_{\text{dark}}$  (Ogren, 2000). For example, higher sugar content was linked to faster  $R_{\text{dark}}$  in leaves of *Quercus rubra* (Griffin *et al.*, 2002, Whitehead *et al.*, 2004), as well as in *Spinacia oleracea* and *Alocasia macrorrhiza* (Noguchi *et al.*, 1996). However, sugars were not correlated to  $R_{\text{dark}}$  in two alpine perennials, *Bisorta bistortoides* and *Campanula rotundifolia* (McCutchan & Monson, 2001). Moreover, thermal acclimation often alters temperature-normalized rates of  $R_{\text{dark}}$ , but such changes can occur without concomitant changes in concentrations of soluble sugars (Zaragoza-Castells *et al.*, 2007). Indeed, sugar concentrations are often relatively constant across a range of environmental treatments (Ayub *et al.*, 2011, Martínez-Vilalta *et al.*, 2016), likely as a

consequence of the synthesis and degradation of starch (Hüve *et al.*, 2012). Starch turnover is regulated by a circadian clock (Graf *et al.*, 2010, Graf & Smith, 2011) and increases during the day as photosynthetic assimilates are typically stored as starch in leaves if not immediately used or exported (Pilkington *et al.*, 2015, Smith & Stitt, 2007). Starch is broken down at a constant rate during the night to provide a relatively constant supply of sugars for  $R_{\text{dark}}$  and growth in the absence of photosynthesis, with approximately 90-95% of the starch consumed by dawn (Graf *et al.*, 2010, Pilkington *et al.*, 2015, Smith & Stitt, 2007). If this circadian regulation of starch stabilizes sugar concentrations over a 24-hour cycle, leaf sugars are unlikely to limit  $R_{\text{dark}}$  and contribute to diel variation. Nevertheless,  $R_{\text{dark}}$  may be affected by diel changes in the availability of other substrates of respiration, specifically amino acids, proteins, organic acids and/or lipids (O'Leary *et al.*, 2011, O'Leary *et al.*, 2017). As described in O'Leary *et al.* (2017), such metabolites may drive respiration by supplying intermediates to the TCA cycle, reductant for ATP synthesis via oxidative phosphorylation, and carbon skeletons required for biosynthesis or nitrogen (N) assimilation into amino acids. Thus, substrate-product relationships across respiratory network must be considered when assessing the interactive effect of growth temperature and time of day on leaf  $R_{\text{dark}}$  and to identify key metabolites that correlate strongly with  $R_{\text{dark}}$ .

Metabolomics may provide insights into factors underpinning variability in fluxes through the respiratory pathways, and thus how  $R_{\text{dark}}$  varies with temperature and diel cycle. One interesting observation from metabolomic studies published to-date is the amount to which metabolite concentrations vary with temperature. Sicher (2015) showed that 28 of 43 total metabolites in soybean leaflets were affected by an increase in growth temperature. With the exception of raffinose and  $\gamma$ -aminobutyric acid (GABA), most of the organic and amino acid metabolites decreased in response to elevated growth temperature. In wheat leaves under warmer night temperatures, the abundance of fumarate and alanine increased, while glutamine, glutamate and GABA remained constant (Impa *et al.*, 2019). Glaubitz *et al.* (2015) found that warmer day (30°C) and night (28°C) temperatures increased the abundance of amino and organic acids in temperature-sensitive rice cultivars (e.g. DR2 and M202) but not in intermediate (e.g. IR64 and IRRI123) and temperature-tolerant cultivars (e.g. IR72 and Taipei309). In addition to being affected by growth temperature, amino acids change in response to the diel cycle. For example, Urbanczyk-Wochniak *et al.* (2005) found amino acid levels increased in the light and decreased in the dark in potato (*Solanum tuberosum*) leaves. The metabolites mentioned above are regulated by the TCA cycle which is central to respiration,

suggesting that the strong response of many metabolites to temperature and light are likely to be associated with alterations in respiratory rates.

TCA cycle intermediates can also respond to changes in growth temperature and the time of day. There is evidence that malate and fumarate accumulate during the day, while citrate, aconitate and succinate accumulate at night (Gibon *et al.*, 2006, Watanabe *et al.*, 2014). Further, light suppresses the pyruvate dehydrogenase complex (PDC) (Tcherkez *et al.*, 2009), potentially reducing PDC activity and its decarboxylation of pyruvate, reducing entry of carbon into the TCA cycle (Tovar-Mendez *et al.*, 2003) with consequences for the concentration of TCA cycle intermediates. Moreover, in some cases the TCA cycle can be supplemented through the action of light-induced phosphoenolpyruvate carboxylate (PEPC) that produces oxaloacetate and malate (Araújo *et al.*, 2012, Marsh *et al.*, 2003, Sullivan *et al.*, 2004). In this pathway, PEPC generally replenishes the TCA cycle intermediates that are depleted when  $\alpha$ -ketoglutarate is used for amino acid synthesis. In addition to these day/night effects, measuring temperature can also alter the abundance of TCA cycle intermediates. For example, the concentrations of  $\alpha$ -ketoglutarate, fumarate, malate, and citrate increased when *Arabidopsis* leaves were cooled from 20°C to 4°C (Cook *et al.*, 2004). In *Arabidopsis* grown at 15°C for 24 hours relative to 20°C controls, succinate, fumarate, and malate accumulated at the beginning of the day but otherwise fluctuated similarly to the 20°C controls during the light period (Florian *et al.* (2014). Further, metabolites that typically vary diurnally – such as alanine, phenylalanine and glutamine – maintain their diurnal oscillations even when leaves are chilled at 4°C (Espinoza *et al.*, 2010). The aforementioned studies imply that there are distinct respiratory metabolite adjustments to temperature and to light/dark cycles. However, more in-depth experimentation of the interaction between temperature and the diel cycle is lacking and should be explored

In the current study, we investigated whether thermal acclimation of respiratory metabolism interacts with the day/night cycle in rice (*Oryza sativa*, cultivar IR64). Rice is one of the most important food crops yet yields are declining, likely from rising growth temperatures (Wassmann *et al.*, 2009). In a recent study, we assessed the effect of growth temperature on acclimation of photosynthesis and respiration in the IR64 cultivar (Rashid *et al.*, 2020); in that study, growth temperature-dependent changes in leaf carbon exchange were assessed at a single time point during the day. Our current study used metabolite profiling to assess how growth temperature and the diel cycle might influence the concentrations of metabolites in the respiratory network of rice. We explore whether adjustments in the abundance of individual metabolites provide insights into how growth temperature affects

regulatory control of respiratory metabolism. We specifically examined: (1) if rates of leaf  $R_{\text{dark}}$  (measured as  $\text{O}_2$  uptake in darkness) vary over a day, (2) whether the extent of the diel variation is influenced by growth temperature, (3) if diel variations in  $R_{\text{dark}}$  are linked to concomitant changes in substrate availability, and (4) the extent to which growth temperature and day/night cycles influence the metabolome profile.

## Material and Methods

### *Plant material and temperature treatments*

Rice (*Oryza sativa* L.) cultivar IR64 was grown hydroponically in a glasshouse facility at the Australian National University in Canberra, Australia. Growth conditions were the same as those reported in Rashid *et al.* (2020). The hydroponic solution consisted of  $\text{NH}_4\text{NO}_3$  (1.4 mM),  $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$  (0.6 mM),  $\text{K}_2\text{SO}_4$  (0.5 mM),  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$  (0.2 mM),  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  (0.8 mM) and micronutrients: Fe-EDTA (0.07 mM),  $\text{H}_3\text{BO}_3$  (0.037 mM),  $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$  (0.009 mM),  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$  (0.00075 mM),  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  (0.0003 mM),  $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$  (0.0001 mM),  $\text{NH}_4\text{VO}_3$  (0.000138 mM) and  $\text{Na}_2\text{SiO}_3$  (0.0012963 mM). Twelve 20 l hydroponic tanks holding a maximum 20 plants were placed in a temperature-controlled glasshouse at 30°C during the day and 25°C (30/25°C) at night under a natural photoperiod. The sunset and sunrise times were 19:30 and 06:30  $\pm$  15 min, respectively. Photosynthetically active radiation (PAR) measured between 10:00 and 15:00 hours ranged from 400 and 1200  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ . After two weeks at 30/25°C, eight tanks were randomly chosen and shifted to adjacent glasshouse rooms set to 25/20°C or 40/35°C temperature regimes (four tanks in each room). Another four tanks were maintained in the original 30/25°C room as controls. Each plant within a separate tank was considered as a biological replicate (Supporting Information Figure S1). Sampling for all experiments unless stated otherwise occurred over a single 24-hour period at 04:30, 08:30, 13:00, 17:30 and 21:30 hours. To minimise developmental differences, all leaf samples for respiratory and metabolite profiling were taken from young fully expanded leaves on days seven and eight after temperature transfer from plants previously unsampled which developed under the prevailing temperature regime.

### *Dark respiration measurements*

Rates of dark respiration ( $R_{\text{dark}}$ ) were measured using a high-throughput fluorophore  $\text{O}_2$ -sensor system (Astec Global, Maarssen, The Netherlands) as previously described (Coast *et al.*, 2019, O'Leary *et al.*, 2017, Scafaro *et al.*, 2017). Four independent plants, each from a separate tank,

from each growth temperature treatment were placed in a container with a small volume of hydroponic solution and brought back to the laboratory to be processed. After 30 min in darkness two separate 2 cm<sup>2</sup> sections from the middle part of the lamina were cut and placed in separate 2 ml capacity O<sub>2</sub>-sensor sampling tubes. One sample tube was measured at a common temperature of 30°C, and the other at the leaf at the prevailing growth temperature.  $R_{\text{dark}}$  was calculated from an O<sub>2</sub> consumption slope between 30 min to two hours after initiation of the fluorophore O<sub>2</sub>-sensor system. At the cessation of measurements, leaf sections were collected, stored in paper bags, oven-dried at 70°C and weighed to obtain dry mass. The dried leaf materials were subsequently used in the determination of starch and soluble sugar concentrations as described below. For expediency of sample collection and processing we did not measure leaf area. For metabolite profiling over the diurnal cycle, leaves from the same plants chosen for respiration were harvested in the glasshouse, immediately frozen in liquid N<sub>2</sub> and stored at minus 80°C.

#### *Determination of soluble sugars, starch and nitrogen concentrations*

To determine the concentrations of soluble sugars and starch, dried rice leaf sections were ground to a fine powder and 5-10 mg placed in a 2 ml microfuge tube. 0.5 ml of 80% (v/v) ethanol was added and vigorously vortexed for 20 sec. The tissues were incubated at 80°C while being orbitally shaken at 500 rpm for 20 min on a Thermomixer (Eppendorf). The samples were then centrifuged for 5 min at 12,000 rpm, and the resulting supernatant and pellet were separated. This procedure was repeated twice again on the same pellet, and supernatant pooled. The supernatant and pellet were used for determination of the concentrations of soluble sugars and starch using a Fructose Assay Kit (Sigma-Aldrich) and a Total Starch Assay Kit (Megazyme), respectively, following manufacturer's instructions. A standard curve for soluble sugars was generated using a series of known concentrations of sucrose, glucose and fructose (Sigma-Aldrich). Measurements were collected using a microtitre plate reader (Infinite M1000Pro; Tecan).

To determine the concentration of nitrogen (N), approximately 1-2 mg dried rice leaf sections were ground and placed in a tin-made capsule. The analysis was conducted using combustion, combining an elemental analyser 15 (Heraeus CHN-O Rapid), a Finnigan MAT Trapping box HT and a Finnigan MAT mass spectrometer (delta D) with a dual inlet at a precision of 0.1 ‰. N concentrations on leaf dry mass basis were calculated based on Gebauer and Schulze (1991).



### *Gas Chromatography-Mass Spectrometry (GC-MS) metabolite analysis*

To quantify metabolites involved in carbon metabolism a mass spectrometry approach using GC-MS metabolite profiling was adopted. The extraction was carried out according to the procedure previously described in Che-Othman *et al.* (2020) with some modifications. Frozen leaf tissue was ground to powder and approximately 25 mg transferred into frozen 2 ml microfuge tubes. 0.5 ml of cold extraction buffer was added, and incubated in a Thermomixer for 20 min at 75°C and 1400 rpm mixing. All tubes were then centrifuged at 20,000  $\times g$  for 10 min at room temperature. 60  $\mu$ l of supernatant was dried down in 200  $\mu$ l glass insert for derivatization prior to metabolite profiling using Gas Chromatography-Mass Spectrometry (GC-MS). The derivatisation was performed using the MPS2 XL-Twister autosampler, Gerstel GmbH & Co. KG). 1  $\mu$ l of derivatised sample was immediately vaporised in the inlet at 250°C and injected onto the chromatography column in split-less mode. Helium was used as carrier gas. Compounds were eluted by the following temperature gradient: hold for 1 min at 70°C then ramp with 7°C/min to 325°C and hold for 3.5 min. The ion transfer line was heated to 280°C and the ion source and quadrupole were at 150°C and 230°C respectively. The resulting peaks were analysed using MassHunter Workstation Software Quantitative Analysis Version B.07.1 / Build 7.1.524.0 for GCMS (Agilent Technologies). The integrated area of the quantifier ion for each peak was compared between samples after normalisation. Metabolites were normalised against the weighted and averaged signal of 3 internal standards followed by weighting against the average measured signal across all samples for each compound before statistical analysis was performed. Metabolite data is presented in Supporting Information dataset S1.

### *Statistical Analyses*

Data were analysed using one-way analyses of variance (ANOVA), two-way ANOVA, principal component analysis (PCA), weighted correlation network analysis, permutational multivariate analysis of variance and multiple comparison tests using the SigmaPlot v11.0 software (Systat Software Inc) and/or R v3.4.4 software (R Core Team, 2018) with RStudio. In R v3.4.4 software, data preparation and plotting were done with packages tidyverse (Wickham, 2017). The PCA was carried out with R packages ‘FactoMineR’ (Le *et al.*, 2008), factoextra (Kassambara & Mundt, 2017) and ade4 (Dray & Dufour, 2007). Metabolite levels were centred and scaled with standard score. Significance testing was done with R package ‘vegan’ (Oksanen *et al.*, 2018). Weighted correlation network analysis was conducted using R package ‘WGCNA’ (Langfelder & Horvath, 2008). Permutational multivariate analysis of

variance was carried out using Euclidean distance matrices of metabolites compared pairwise (Anderson, 2001, McArdle & Anderson, 2001). The analysis produces test statistic analogous to Fisher's F-ratio, and calculates P-value through permutations.

## Results

### *R<sub>dark</sub> varies independently with time of day and temperature*

When measured at the prevailing growth temperature of each treatment,  $R_{\text{dark}}$  varied with time of day (Fig. 1a), with rates typically being at their lowest in the early evening (noting that night temperatures were 5°C lower than those during the day).  $R_{\text{dark}}$  was faster in the hottest treatment (40/35°C) and slower at the two lower growth temperatures (warm, 30/25°C; cool, 25/20°C), and this effect was consistent through the day with no significant growth temperature time interaction (Fig. 1a, Supporting Information Table S1).

Despite prevailing rates of  $R_{\text{dark}}$  rising with growth temperature (Fig. 1a), there was evidence that acclimation did occur, as the hot-grown plants exhibited lower temperature-normalized rates of  $R_{\text{dark}}$  (i.e. lower rates measured at 30°C) than their warm- and cool-grown counterparts (Fig. 1b, Supporting Information Table S1). This finding supported our previous findings of respiratory thermal acclimation in rice, quantified via measurements of respiratory CO<sub>2</sub> release in darkness during the daylight period of 10:00-14:00 hours (Rashid *et al.*, 2020). The acclimation response (determined as the difference among the three growth temperatures in temperature-normalized rates of leaf  $R_{\text{dark}}$ ) appeared greater at 17:30 hours (2 h before sunset) and lesser during the night (Fig. 1b). However, the interaction between growth temperature and time of day was not statistically significant ( $P=0.065$ ), suggesting that the effect of growth temperature on  $R_{\text{dark}}$  is relatively similar through the diel cycle.

### *Diel variations in R<sub>dark</sub> are not linked to concomitant changes in sugar substrate availability*

Changes in starch and soluble sugar content (Fig. 2) did not mirror the acclimation of – or the diel variation in – rates of  $R_{\text{dark}}$  measured at 30°C (Fig. 1b). The concentration of individual and total leaf soluble sugars remained constant throughout the day (Figs 2a-d), with no particular sugar exhibiting a statistically significant response to time of day. Total soluble sugar concentration was affected by growth temperature, being lower in the cold (25/20°C), mainly due to lower glucose and fructose levels in leaves grown in colder temperatures (Figs 2a-d). Unlike soluble sugars, starch concentration varied diurnally irrespective of growth temperature, decreasing through the night and increasing through the day, with a minimum value near

sunrise and maximum value near sunset (Fig. 2e). Low growth temperature resulted in significantly greater leaf N concentrations compared to warm- and hot-grown plants (Fig. 2f); while there was a significant main effect of time on leaf N, there was no clear diel pattern.

#### *Metabolite profiles: relationships among metabolites*

Two general findings were observed from metabolite profiling. Firstly, principal component analysis (PCA) of correlations among metabolites (Fig. 3a) revealed a negative correlation between amino acids and carbohydrates/organic acids. Secondly, changes in metabolites in response to a day/night cycle or growth temperature were mostly independent of one another (Figs 3b,c, Table 1). Independence was evident from metabolite responses to day and night being separated mostly along the first principal component. In contrast, metabolite responses to growth temperature were grouped mostly along the second principal component (Fig. 3b,c). Apart from the shikimate pathway derived tryptophan and tyrosine, a two-way ANOVA corroborated that there was little significant interaction among metabolites responding to changes in day/night cycle and growth temperature (Table 1).

#### *Metabolite profiles: The influence of growth temperature and day/night cycles*

Since there was minimal interaction between the diurnal cycle and growth temperature in terms of metabolite adjustments, we performed separate one-way ANOVA, one for the diel cycle, and the other for growth temperature changes in metabolites. We displayed these changes in relation to the respiratory pathway (Figs 4 and 5). Generally, the diel changes in metabolites and metabolite responses to growth temperature did not align with specific metabolic pathways (i.e. glycolysis, TCA cycle, shikimate pathway, amino acid synthesis). For example, a change in the TCA cycle intermediate, aconitate, was not preceded by a change in the antecedent metabolite citrate, nor were changes in aconitate followed by the changes in subsequent metabolites succinate, fumarate, and malate. The most striking general diel and temperature driven change in metabolites was that many branched chain and aromatic amino acids increased in abundance during the night and in response to warmer temperatures (Figs 4 and 5). By contrast, short chained amino acids such as alanine, glycine and serine, as well as the N side chain amino acids glutamine and asparagine, were in greater abundance during the day and at cooler growth temperatures.

Independent of the growth temperature, most of the TCA cycle intermediates changed with diel cycle (Fig. 4). The intermediate substrates at the beginning of the glycolysis pathway (glucose-6-phosphate and fructose-6-phosphate) varied significantly through the 24-hour cycle,

with abundances highest at mid-day and lowest at night (Fig. 4). Citrate accumulated at night,  $\alpha$ -ketoglutarate accumulated during the day, and malate depleted during the day. The amino acid alanine was more abundant during the day and depleted at night, but aspartic acid was reduced in the day and more abundant at night. Tryptophan, phenylalanine and tyrosine, aromatic amino acid products of the shikimate pathway, were significantly reduced at night (Fig. 4).

In contrast to diel cycle, growth temperature alone affected fewer metabolites (Fig. 5). This may be a consequence of the diel cycle having both an irradiance and temperature component (with nights being 5°C cooler than that during the day). The metabolite analyses revealed that growth temperature treatment subtly changed relative abundance of soluble sugars, confirming the enzymatic assay of total soluble sugars (Fig. 2). Cold temperature reduced abundance of glucose and fructose, albeit not to a significant level. By contrast, increasing growth temperature greatly reduced levels of glucose-6-phosphate and fructose-6-phosphate. As with diel cycle, phenylalanine and tyrosine were significantly affected by temperature with lower abundance in the colder grown rice. Intermediates of the TCA cycle did not significantly vary with growth temperature apart from aconitate which increased with increasing growth temperature. Many amino acid pools generated from TCA intermediates significantly changed in response to growth temperature, with lysine and GABA increasing at the hottest temperature of 40/35°C, proline and isoleucine decreasing at the lowest temperature of 25/20°C, and glutamine increasing at 25/20°C (Fig. 5).

#### *Metabolite profiles: relationships with $R_{\text{dark}}$*

Weighted correlation network analysis identified seven metabolites that correlated with changes in  $R_{\text{dark}}$  measured at a common 30°C (Fig. 6, Supporting Information Table S3). Correlations between  $R_{\text{dark}}$  and metabolites were observed with changes in growth temperature, the diel period, or both. Weighted correlation network analysis describes correlation patterns in data (Zhang & Horvath, 2005) and has been successfully used to analyse biological data such as gene co-expression (Stuart *et al.*, 2003, Weston *et al.*, 2008) and protein-protein interactions (Jeong *et al.*, 2001). Applied here, it described the interactions among metabolites by grouping similarity responding metabolites into clusters and identifying seven key metabolites with responses representative of the clusters. This allowed for a test of relationship between the key metabolites and  $R_{\text{dark}}$  at 30°C (Fig. 6), without correcting for multiple comparisons necessary if all metabolites were tested. Metabolites xylitol, aconitate, and erythritol increased with declining  $R_{\text{dark}}$  at 30°C, primarily in association with growth

temperature (Fig. 6). Serine, putrescine, and fructose-6-phosphate increased with rising  $R_{\text{dark}}$ , associated with both temperature and diel responses. Succinate increased with rising  $R_{\text{dark}}$  primarily associated with diel responses alone (Fig. 6).

## Discussion

Our study has pointed to a surprising role of TCA-dependent metabolites – particularly amino acids – in the acclimation response of respiration. Glycolysis and the TCA cycle derived xylitol, erythritol, serine, and putrescine were strongly correlated (although not necessarily due to a causal relationship) with adjustments in  $R_{\text{dark}}$  to growth temperature and the day/night cycle (Fig. 6). More expectedly, some of the intermediates of glycolysis (fructose-6-phosphate) and the TCA cycle (aconitate and succinate) correlated with adjustments in  $R_{\text{dark}}$ , although others (malate and citrate) did not (Fig. 6). Many of the metabolites that did change in abundance to growth temperature or the diel cycle did so for one and not the other (Table 1), supporting separate influences of growth temperature and diel perturbations on respiration. The limited interaction between metabolite responses to growth temperature and a day/night cycle was evident in respiratory flux measurements, with the acclimation phenotype largely been held across the diurnal cycle (Fig. 1b). However, of interest, the shikimate derived aromatic amino acids were the only metabolites that significantly adjusted in relative abundance to both growth temperature and the diel period. These observations provide insights into the likely metabolic determinants of respiratory acclimation and its response to the day/night cycle.

### *Does time of day influence our conclusion of whether acclimation has occurred?*

While temperature-normalized rates of  $R_{\text{dark}}$  of rice leaves differed among the growth temperature treatments (indicating that thermal acclimation occurred) and varied through the diurnal cycle, the effect of growth temperature did not vary significantly with time (Fig. 1b, Supporting Information Table S1). Thus, it appears that – at least for rice used in our study – respiratory thermal acclimation can be assessed using measurements made at a single time point during the day. Expressing  $R_{\text{dark}}$  on a leaf area basis is unlikely to alter the result as leaf mass per area did not change in rice exposed to the same experimental conditions (Rashid *et al.* 2020). As noted in the Introduction, temperature-normalized rates of  $R_{\text{dark}}$  – measured at a single time point during the day – are commonly used to quantify thermal acclimation and biogeographical patterns in leaf respiration (Atkin *et al.*, 2005a, Huntingford *et al.*, 2017, Reich *et al.*, 2016, Slot & Kitajima, 2015, Smith & Dukes, 2013). Such studies point to decreases in

temperature-normalized  $R_{\text{dark}}$  with sustained exposure to higher growth temperature, and lower rates of temperature-normalized  $R_{\text{dark}}$  in the plants growing in the tropics than the arctic (Atkin et al. 2015). Our results suggest that those conclusions are unlikely to be affected by the time of day when rates of leaf temperature-normalized  $R_{\text{dark}}$  are measured.

#### *Specific metabolites that correlate with $R_{\text{dark}}$*

Changes due to growth temperature in the sugar alcohols xylitol and erythritol, and the TCA organic acid aconitate, had the strongest correlation with  $R_{\text{dark}}$  at 30°C (Fig. 4). In a recent study, warmer night temperatures led to an increase in the sugar alcohol myo-inositol in spikes of a relatively heat tolerant wheat cultivar (Impa et al., 2019). The authors attributed this to the role of myo-inositol to osmolyte protection. The increase in xylitol and erythritol that we observed is likely to be a similar osmolyte protection response to heat. Increases in these sugar alcohols and reduced  $R_{\text{dark}}$  at 30°C – while inversely correlated - may be independent responses to increased  $T$ . Alternatively, the accumulation of osmolytes for heat protection may in some way suppress respiratory requirements of the leaf. Stabilisation of membranes as a result of osmolyte protection during heat acclimation is postulated (Gauthier et al., 2014, Sung et al., 2003). As such, a greater maintenance of proton motive force, through reduced membrane leakiness, is a conceivable way in which respiratory costs could be reduced with increases in xylitol and erythritol.

While growth temperature had no significant effect on succinate levels (Fig. 5), time of day did significantly alter the abundance of this TCA intermediate, with succinate concentrations declining through the night (Fig. 4). When succinate abundance was plotted against  $R_{\text{dark}}$  at 30°C, (Fig. 6), a significant positive correlation was observed, suggesting that succinate may be a key control point in coordinating the TCA cycle with respiratory adjustments over a day/night cycle. This is likely associated with the light dependent regulation of succinate dehydrogenase activity and its influence over succinate content (Daloso et al., 2015, Eprintsev et al., 2013).

The amino acid serine, the amino acid breakdown product putrescine, and the phosphorylated sugar fructose-6-phosphate were the metabolites with strongest correlation between  $R_{\text{dark}}$  at 30°C associated with both growth temperature and the diel cycle (Fig. 6). It is interesting that an amino acid and its associated amino acid breakdown product correlate with  $R_{\text{dark}}$  as strongly as a sugar substrate of glycolysis. The correlation between fructose-6-phosphate and  $R_{\text{dark}}$  at 30°C, both over the diel cycle, and growth temperature, is understandable as this initial substrate of glycolysis is a regulatory point of respiration (Plaxton

& Podestá, 2006, Stitt, 1990). Serine and putrescine, on the other hand, are not intermediates of the TCA cycle. As discussed below, there seems to be a tight coordination between amino acid synthesis and respiration which may account for this close association between serine, putrescine, and the respiratory flux at 30°C. Additionally, serine is a product of photorespiration, which may explain its variability across growth temperatures and the day and night periods (Fig. 6). Photorespiration would not occur in the dark and would be stimulated by heating (Walker *et al.* 2016). Levels of serine were lower at night in agreement with reduced photorespiration. However, serine levels were often lower at 40°C than 25°C in the day (Fig. 6), at odds with photorespiratory expectations. This highlights the difficulty in characterising the relationships between metabolite pools and environmental perturbations. Many metabolites will have interconnected pathways that are influenced separately by temperature and light, such as the involvement of serine in photorespiration and protein synthesis.

#### *Assimilate utilisation*

Starch accumulated during the day in leaves, and was depleted during the night, reaching a minimum in the early morning (Fig. 2e) - a pattern that is well documented (Graf *et al.*, 2010, Graf & Smith, 2011). Interestingly, starch concentrations exhibited a similar pattern irrespective of growth temperature, with no significant interaction between temperature and the starch diurnal cycle. Increased catabolism of sugars did not supplement  $R_{\text{dark}}$  substrate supply as soluble sugar content was not depleted at 40/35°C (Fig. 2a-d). Thus, starch was depleted from a similar starting point and at a similar rate between 30/25°C and 40/35°C grown plants despite the latter having much faster  $R_{\text{dark}}$ .  $^{14}\text{CO}_2$  pulse-chase experiments demonstrate that faster rates of  $R_{\text{dark}}$  are associated with faster depletion of starch (Hüve *et al.*, 2012). However, transport of starch out of leaves, and utilisation of starch for growth, also have a major impact on leaf starch content (Stitt & Zeeman, 2012). Rice that we grew under the same experimental conditions do have faster leaf elongation rates at 40/35°C compared to the cooler temperatures, across the entire diel cycle (Rashid *et al.* 2020). The similar pattern of starch drawdown despite differences in  $R_{\text{dark}}$  is therefore likely due to temperature mediated shifts in starch transport (as sucrose) out of leaves to other organs, and different rates of utilisation due to temperature-dependent differences in leaf biomass accumulation. In other words, the levels of starch and soluble sugars were relatively constant across growth temperatures, but the fluxes of these respiratory substrates likely increased with heat to match stimulated growth.

#### *Amino acids respond strongly to the diurnal cycle and growth temperature*

Although metabolite levels are not equivalent to fluxes, increased abundance of amino acids in the dark seems counterintuitive, as N incorporation into amino acids predominantly occurs in the light (Canvin & Atkins, 1974, Fritz *et al.*, 2006, Nelson *et al.*, 2014). Toxic ammonia is initially incorporated into glutamine and glutamate by glutamine synthetase and glutamate 2-oxoglutarate aminotransferase, an energy expensive process likely supplemented by photosynthetic energy generation (Plaxton & Podestá, 2006, Tcherkez *et al.*, 2017). However, we saw a greater accumulation of certain amino acids at night. In particular, the abundance of longer chained and aromatic amino acids, such as the shikimate pathway derived aromatics phenylalanine, tyrosine and tryptophan; the oxaloacetate derived aspartic acid and lysine; and the  $\alpha$ -ketoglutarate derived proline and GABA, all significantly increased at night or with warmer growth temperature (Fig. 4, 5). Of further interest, this increase in long chained and aromatic amino acids had a strong negative relationship with carbohydrate/organic acid pools (Fig. 3a) This suggests that N was likely fixed into amino acids during the light, but catabolised at night (more so when nights were warmer) through the respiratory pathways to synthesise a wider array of amino acids, at the expense of carbohydrate and organic acid levels. Alanine is the prime candidate for night-time N recycling, as it is more abundant than other amino acids in the day and significantly depleted at night. It also has the shortest anabolic/catabolic pathway, being converted between alanine and pyruvate by alanine aminotransferases (Hildebrandt *et al.*, 2015). Furthermore, the *de novo* synthesis of alanine in the light is faster than other amino acids (Ishihara *et al.*, 2015, Nelson *et al.*, 2014, Szecowka *et al.*, 2013), indicating its role in initial N storage and carbon skeleton synthesis. Glycine and serine are highly abundant, short-chained amino acids that accumulate during the day and are depleted at night (Fritz *et al.*, 2006), likely playing a similar role to alanine in night-time N and carbon recycling. Recycling of organic carbon skeletons through the TCA cycle may coordinate substrate supply with environmental dependent shifts in energy requirements, maintaining balanced redox poise (Igamberdiev & Eprintsev, 2016).

#### *TCA cycle organic acid intermediates*

Neither irradiance nor growth temperature led to a uniform shift in TCA cycle intermediates (Fig. 4, 5, Table 1). This is likely attributable to light-dependent regulation of TCA enzymes. Succinate dehydrogenase and fumarase are downregulated in the light via phytochrome A (Eprintsev *et al.*, 2013, Eprintsev *et al.*, 2016) as well as deactivated by a thioredoxin enzyme (Daloso *et al.*, 2015) leading to malate, fumarate and succinate increasing during high-light,



highly-reduced conditions. This tightly controlled light-dependent regulation of succinate dehydrogenase is the likely reason that diel variations in succinate exhibited the strongest correlation with  $R_{\text{dark}}$  at 30°C (Fig. 4). In addition, the anaplerotic pathway involving the phosphorylation by PEPC (O’Leary *et al.*, 2011, Tcherkez *et al.*, 2009, Werner *et al.*, 2011) could also contribute to the accumulation of malate and fumarate. Aconitate isomerase activity is greater in the day than at night which may explain a decrease in citrate levels during the day (Igamberdiev & Eprintsev, 2016). We can therefore attribute the significant shifts in TCA intermediates like malate, citrate, fumarate and succinate between day and night (Fig. 4) to shifts in enzyme velocities regulated by the day night cycle. Interestingly, the limited effect of growth temperature on succinate, malate or citrate levels (Fig. 5) indicates that these day/night enzyme activity dependent fluctuations are not disrupted by longer-term acclimation to growth temperature.

The accumulation of malate during the day and its depletion at night is linked to enzymatic regulation, with evidence that NAD-malic enzyme is inactive in the light but not dark, coordinating malate storage and utilization with the day/night cycle (Hill, 1992). Additionally, a link between photosynthesis, malate concentrations and leaf respiration is evident in Arabidopsis where reduced plastid NAD-malate dehydrogenase activity results in higher night-time malate and starch content and reduced  $R_{\text{dark}}$  (Beeler *et al.*, 2014). Similarly, malate depletion due to overexpression of NADP-malate dehydrogenase in Arabidopsis chloroplasts leads to a suppression of  $R_{\text{dark}}$  and ultimately plant starvation (Zell *et al.*, 2010). Our observation of malate declining at night can thus be attributed to malate synthesis during the day as a storage of photosynthesis reducing potential, followed by oxidation at night to facilitate respiration. The use of stored malate as a TCA substrate at night would likely be another contributor to the disparity between growth temperature dependent  $R_{\text{dark}}$  and starch drawdown we observed.

### Conclusions

From this work we have shown that  $R_{\text{dark}}$ , glycolysis, and the TCA cycle are highly responsive to both the day/night cycle and temperature acclimation. However, there is little interaction, with minimal influence of the diel cycle on temperature acclimation. Shikimate pathway derived aromatic amino acids were the only metabolites to interact in response to both growth temperature and the day/night cycle. The glycolysis substrate fructose-6-phosphate and the TCA cycle intermediates aconitate and succinate correlate strongly with environmental driven acclimation in  $R_{\text{dark}}$ . Thus, there is evidence of the tight coordination of glycolysis and the TCA

cycle to respiratory acclimation. However, the relationship between  $R_{\text{dark}}$  acclimation and the metabolites xylitol, erythritol, putrescine, and serine suggests that by-products of the respiratory pathways are similarly associated with  $R_{\text{dark}}$  acclimation to temperature and photoperiod. Our results point to a coordinated and complex regulatory system, whereby environmental shifts in redox potential and ATP demand are balanced through adjustments in anaplerotic reactions of the TCA cycle and consequential changes in  $R_{\text{dark}}$ . How these metabolite adjustments ultimately influence plant growth, development and reproductive fitness may have significant implications on plant adaptation to a warmer climate and crop yields.

### Acknowledgements

This work was supported by the Australian Research Council (ARC) Centre of Excellence in Plant Energy Biology (CE140100008). F.A.A.R was supported by the Ministry of Education, Malaysia.

### Author Contribution

FAAR, APS, RCD, JM, NLT, and OKA conceptualised experiments. FAAR, APS, SA, and RF performed experimentation and/or data analysis. FAAR, APS, SA, NLT and OKA drafted the manuscript with contributions from all authors.

### References

- Amthor JS. 2000.** The McCree–de Wit–Penning de Vries–Thornley respiration paradigms: years later. *Annals of Botany*, **86**, 1-20.
- Anderson MJ. 2001.** A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, **26**, 32-46.
- Araújo WL, Nunes-Nesi A, Nikoloski Z, Sweetlove LJ, Fernie AR. 2012.** Metabolic control and regulation of the tricarboxylic acid cycle in photosynthetic and heterotrophic plant tissues. *Plant, Cell & Environment*, **35**, 1-21.
- Atkin OK, Bloomfield KJ, Reich PB, Tjoelker M, Asner GP, Bonal D, Bönisch G, Bradford MG, Cernusak LA, Cosio EG et al. 2015.** Global variability in leaf

- respiration in relation to climate, plant functional types and leaf traits. *New Phytologist*, **206**, 614-636.
- Atkin OK, Bruhn D, Hurry VM, Tjoelker MG. 2005a.** Evans Review No. 2: The hot and the cold: unravelling the variable response of plant respiration to temperature. *Functional Plant Biology*, **32**, 87-105.
- Atkin OK, Bruhn D, Tjoelker MG. 2005b.** Response of plant respiration to changes in temperature: Mechanisms and consequences of variations in  $Q_{10}$  values and acclimation. In: *Plant respiration: From cell to ecosystem*. (eds Lambers H, Ribas-Carbo M) pp 95-135. Dordrecht, Netherlands, Springer.
- Atkin OK, Macherel D. 2009.** The crucial role of plant mitochondria in orchestrating drought tolerance. *Annals of Botany*, **103**, 581-597.
- Ayub G, Smith RA, Tissue DT, Atkin OK. 2011.** Impacts of drought on leaf respiration in darkness and light in *Eucalyptus saligna* exposed to industrial-age atmospheric CO<sub>2</sub> and growth temperature. *New Phytologist*, **190**, 1003-1018.
- Beeler S, Liu H-C, Stadler M, Schreier T, Eicke S, Lue W-L, Truernit E, Zeeman SC, Chen J, Kötting O. 2014.** Plastidial NAD-dependent malate dehydrogenase is critical for embryo development and heterotrophic metabolism in *Arabidopsis*. *Plant Physiology*, **164**, 1175.
- Canvin DT, Atkins CA. 1974.** Nitrate, nitrite and ammonia assimilation by leaves: Effect of light, carbon dioxide and oxygen. *Planta*, **116**, 207-224.
- Che-Othman MH, Jacoby RP, Millar AH, Taylor NL. 2020.** Wheat mitochondrial respiration shifts from the tricarboxylic acid cycle to the GABA shunt under salt stress. *New Phytologist*, **225**, 1166-1180.
- Coast O, Shah S, Ivakov A, Gaju O, Wilson PB, Posch BC, Bryant CJ, Negrini ACA, Evans JR, Condon AG. et al. 2019.** Predicting dark respiration rates of wheat leaves from hyperspectral reflectance. *Plant, Cell & Environment*, **42**, 2133-2150.
- Cook D, Fowler S, Fiehn O, Thomashow MF. 2004.** A prominent role for the CBF cold response pathway in configuring the low-temperature metabolome of *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 15243-15248.
- Daloso DM, Müller K, Obata T, Florian A, Tohge T, Bottcher A, Riondet C, Bariat L, Carrari F, Nunes-Nesi A, et al. 2015.** Thioredoxin, a master regulator of the tricarboxylic acid cycle in plant mitochondria. *Proceedings of the National Academy of Sciences*, **112**, E1392.

- Davy R, Esau I, Chernokulsky A, Outten S, Zilitinkevich S. 2017.** Diurnal asymmetry to the observed global warming. *International Journal of Climatology* **37**, 79–93.
- Dray S, Dufour A-B. 2007.** The ade4 Package: Implementing the duality diagram for ecologists. *Journal of Statistical Software*, **22(4)**, 1-20.
- Eprintsev AT, Fedorin DN, Igamberdiev AU. 2013.** Ca<sup>2+</sup> is involved in phytochrome A-dependent regulation of the succinate dehydrogenase gene *sdh1-2* in Arabidopsis. *Journal of Plant Physiology*, **170**, 1349-1352.
- Eprintsev AT, Fedorin DN, Sazonova OV, Igamberdiev AU. 2016.** Light inhibition of fumarase in Arabidopsis leaves is phytochrome A-dependent and mediated by calcium. *Plant Physiology and Biochemistry*, **102**, 161-166.
- Espinoza C, Degenkolbe T, Caldana C, Zuther E, Leisse A, Willmitzer L, Hinch, DK, Hannah MA. 2010.** Interaction with diurnal and circadian regulation results in dynamic metabolic and transcriptional changes during cold acclimation in Arabidopsis. *PloS one*, **5**, e14101-e14101.
- Florian A, Nikoloski Z, Sulpice R, Timm S, Araújo WL, Tohge T, Bauwe H, Fernie AR. 2014.** Analysis of short-term metabolic alterations in Arabidopsis following changes in the prevailing environmental conditions. *Molecular Plant*, **7**, 893-911.
- Fritz C, Palacios-Rojas N, Feil R, Stitt M. 2006.** Regulation of secondary metabolism by the carbon–nitrogen status in tobacco: nitrate inhibits large sectors of phenylpropanoid metabolism. *The Plant Journal*, **46**, 533-548.
- Gauthier PPG, Crous KY, Ayub G, Duan H, Weerasinghe LK, Ellsworth DS, Tjoelker MG, Evans JR, Tissue DT, Atkin OK. 2014.** Drought increases heat tolerance of leaf respiration in Eucalyptus globulus saplings grown under both ambient and elevated atmospheric [CO<sub>2</sub>] and temperature. *Journal of Experimental Botany*, **65**, 6471-6485.
- Gebauer G, Schulze E-D. 1991.** Carbon and nitrogen isotope ratios in different compartments of a healthy and a declining Picea abies forest in the Fichtelgebirge, NE Bavaria. *Oecologia*, **87**, 198-207.
- Gibon Y, Pyl E-T, Sulpice R, Lunn JE, Höhne M, Günther M, Stitt M. 2009.** Adjustment of growth, starch turnover, protein content and central metabolism to a decrease of the carbon supply when Arabidopsis is grown in very short photoperiods. *Plant, Cell & Environment*, **32**, 859-874.
- Gibon Y, Usadel B, Blaessing OE, Kamlage B, Hoehne M, Trethewey R, Stitt M. 2006.** Integration of metabolite with transcript and enzyme activity profiling during diurnal cycles in Arabidopsis rosettes. *Genome Biol*, **7**, R76.

- Glaubitz U, Erban A, Kopka J, Hinch DK, Zuther E. 2015.** High night temperature strongly impacts TCA cycle, amino acid and polyamine biosynthetic pathways in rice in a sensitivity-dependent manner. *Journal of Experimental Botany*, **66**, 6385-6397.
- Gonzalez-Meler MA, Taneva L, Trueman RJ. 2004.** Plant respiration and elevated atmospheric CO<sub>2</sub> concentration: Cellular responses and global significance. *Annals of Botany*, **94**, 647-656.
- Graf A, Schlereth A, Stitt M, Smith AM. 2010.** Circadian control of carbohydrate availability for growth in Arabidopsis plants at night. *Proceedings of the National Academy of Sciences*, **107**, 9458-9463.
- Graf A, Smith AM. 2011.** Starch and the clock: the dark side of plant productivity. *Trends in Plant Science*, **16**, 169-175.
- Griffin KL, Turnbull M, Murthy R. 2002.** Canopy position affects the temperature response of leaf respiration in *Populus deltoides*. *New Phytologist*, **154**, 609-619.
- Hildebrandt Tatjana m, Nunes nesi A, Araújo Wagner I, Braun H-P. 2015.** Amino acid catabolism in plants. *Molecular Plant*, **8**, 1563-1579.
- Hill SA. 1992.** Malate metabolism and light-enhanced dark respiration in barley mesophyll protoplasts. In: *Molecular, biochemical and physiological aspects of plant respiration*. (eds Lambers H, Van Der Plas LHW) pp 221-230. The Hague, The Netherlands, SPB Academic Publishing bv.
- Huntingford C, Atkin OK, Martinez-de la Torre A, Mercado LM, Heskell MA, Harper AB, Bloomfield KJ, O'Sullivan OS, Reich PB, Wythers KR *et al.* 2017.** Implications of improved representations of plant respiration in a changing climate. *Nature Communications*, **8**, 1602.
- Hurry V, Igamberdiev AU, Keerberg O, Pärnik T, Atkin OK, Zaragoza-Castells J, Gardenström P. 2005.** Respiration in photosynthetic cells: gas exchange components, interactions with photorespiration and the operation of mitochondria in the light. In: *Plant respiration: from cell to ecosystem*. (eds Lambers H, Ribas-Carbó M) pp 43-61. Dordrecht, the Netherlands, Springer.
- Hüve K, Bichele I, Ivanova H, Keerberg O, Pärnik T, Rasulov B, Tobias M, Niinemets Ü. 2012.** Temperature responses of dark respiration in relation to leaf sugar concentration. *Physiologia Plantarum*, **144**, 320-334.
- Igamberdiev AU, Eprintsev AT. 2016.** Organic Acids: The pools of fixed carbon involved in redox regulation and energy balance in higher plants. *Frontiers in Plant Science*, **7**, 1042-1042.

- Impa SM, Sunoj VSJ, Krassovskaya I, Bheemanahalli R, Obata T, Jagadish SVK. 2019.** Carbon balance and source-sink metabolic changes in winter wheat exposed to high night-time temperature. *Plant, Cell & Environment*, **42**, 1233-1246.
- Ishihara H, Obata T, Sulpice R, Fernie AR, Stitt M. 2015.** Quantifying protein synthesis and degradation in Arabidopsis by dynamic  $^{13}\text{CO}_2$  labeling and analysis of enrichment in individual amino acids in their free pools and in protein. *Plant Physiology*, **168**, 74.
- Jeong H, Mason SP, Barabási AL, Oltvai ZN. 2001.** Lethality and centrality in protein networks. *Nature*, **411**, 41-42.
- Kassambara A, Mundt F. 2017.** factoextra: Extract and visualize the results of multivariate data analyses. R package version 1.0.5.
- Kromer S. 1995.** Respiration during photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, **46**, 45-70.
- Lambers H, Ribas-Carbo M. 2005.** *Plant respiration: From cell to ecosystem*, Dordrecht, Netherlands, Springer.
- Langfelder P, Horvath S. 2008.** WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics*, **9**.
- Le S, Josse J, Husson F. 2008.** FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software* **25**, 1-18.
- Marsh JT, Sullivan S, Hartwell J, Nimmo HG. 2003.** Structure and expression of phosphoenolpyruvate carboxylase kinase genes in solanaceae. A novel gene exhibits alternative splicing. *Plant Physiology*, **133**, 2021-2028.
- Martínez-Vilalta J, Sala A, Asensio D, Galiano L, Hoch G, Palacio S, Piper FI, Lloret F. 2016.** Dynamics of non-structural carbohydrates in terrestrial plants: a global synthesis. *Ecological Monographs*, **86**, 495-516.
- McArdle BH, Anderson MJ. 2001.** Fitting multivariate models to community data: A comment on distance-based redundancy analysis. *Ecology*, **82**, 290-297.
- McCutchan CL, Monson RK. 2001.** Night-time respiration rate and leaf carbohydrate concentrations are not coupled in two alpine perennial species. *New Phytologist*, **149**, 419-430.
- Nelson CJ, Alexova R, Jacoby RP, Millar AH. 2014.** Proteins with high turnover rate in Barley leaves estimated by proteome analysis combined with in planta isotope labeling. *Plant Physiology*, **166**, 91-108.

- Noguchi K, Sonoike K, Terashima I. 1996.** Acclimation of respiratory properties of leaves of *Spinacia oleracea* L., a sun species, and of *Alocasia macrorrhiza* (L.) G. Don., a shade species, to changes in growth irradiance. *Plant and Cell Physiology*, **37**, 377-384.
- Noguchi K, Yoshida K. 2008.** Interaction between photosynthesis and respiration in illuminated leaves. *Mitochondrion*, **8**, 87-99.
- O'Leary B, Park J, Plaxton WC, 2011.** The remarkable diversity of plant PEPC (phosphoenolpyruvate carboxylase): recent insights into the physiological functions and post-translational controls of non-photosynthetic PEPCs. *Biochem J*, **436**, 15-34.
- O'Leary BM, Lee CP, Atkin OK, Cheng R, Brown TB, Millar AH. 2017.** Variation in leaf respiration rates at night correlates with carbohydrate and amino acid supply. *Plant Physiology*, **174**, 2261.
- Ogren E. 2000.** Maintenance respiration correlates with sugar but not nitrogen concentration in dormant plants. *Physiologia Plantarum*, **108**, 295-299.
- Oksanen J, Blanchet FG, Friendly M, et al. (2018)** vegan: Community Ecology Package. R package version 2.5-2.
- Peng S, Huang J, Sheehy JE, Laza RC, Visperas RM, Zhong X, Centeno GS, Khush GS, Cassman KG. 2004.** Rice yields decline with higher night temperature from global warming. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 9971-9975.
- Pilkington SM, Encke B, Krohn N, Hohne M, Stitt M, Pyl ET. 2015.** Relationship between starch degradation and carbon demand for maintenance and growth in *Arabidopsis thaliana* in different irradiance and temperature regimes. *Plant Cell Environ*, **38**, 157-171.
- Plaxton WC, Podestá FE. 2006.** The functional organization and control of plant respiration. *Critical Reviews in Plant Sciences*, **25**, 159-198.
- R Core Team. 2018.** A language and environment for statistical computing. R Foundation for Statistical Computing.
- Rashid FAA, Crisp PA, Zhang Y, Berkowitz O, Pogson BJ, Day DA, Masle J, Dewar RC, Whelan J, Atkin OK, Scafaro AP. 2020.** Molecular and physiological responses during thermal acclimation of leaf photosynthesis and respiration in rice. *Plant Cell & Environment*, **43**, 594-610.
- Reich PB, Sendall KM, Stefanski A, Wei X, Rich RL, Montgomery RA. 2016.** Boreal and temperate trees show strong acclimation of respiration to warming. *Nature*, **531**, 633-636.

- Scafaro AP, Negrini ACA, O’Leary B, Rashid FAA, Hayes L, Fan Y, Zhang Y, Chochois V, Badger MR, Millar AH, Atkin OK. 2017.** The combination of gas-phase fluorophore technology and automation to enable high-throughput analysis of plant respiration. *Plant Methods*, **13**, 16.
- Shameer S, Ratcliffe RG, Sweetlove LJ. 2019.** Leaf energy balance requires mitochondrial respiration and export of chloroplast NADPH in the light. *Plant Physiology*, **180**, 1947-1961.
- Sicher RC. 2015.** Temperature shift experiments suggest that metabolic impairment and enhanced rates of photorespiration decrease organic acid levels in Soybean leaflets exposed to supra-optimal growth temperatures. *Metabolites*, **5**, 443-454.
- Slot M, Kitajima K. 2015.** General patterns of acclimation of leaf respiration to elevated temperatures across biomes and plant types. *Oecologia*, **177**, 885-900.
- Smith AM, Stitt M. 2007.** Coordination of carbon supply and plant growth. *Plant, cell & Environment*, **30**, 1126-1149.
- Smith NG, Dukes JS. 2013.** Plant respiration and photosynthesis in global-scale models: incorporating acclimation to temperature and CO<sub>2</sub>. *Global Change Biology*, **19**, 45-63.
- Smith NG, Dukes JS. 2017.** Short-term acclimation to warmer temperatures accelerates leaf carbon exchange processes across plant types. *Global Change Biology*, **23**, 4840-4853.
- Steer BT. 1973.** Diurnal variations in photosynthetic products and nitrogen metabolism in expanding leaves. *Plant Physiology*, **51**, 744-748.
- Stitt M. 1990.** Fructose-2,6-Bisphosphate as a regulatory molecule in plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, **41**, 153-185.
- Stitt M, Zeeman SC. 2012.** Starch turnover: pathways, regulation and role in growth. *Current Opinion in Plant Biology*, **15**, 282-292.
- Stuart JM, Segal E, Koller D, Kim SK. 2013.** A gene-coexpression network for global discovery of conserved genetic modules. *Science*, **302**, 249.
- Sullivan S, Jenkins GI, Nimmo HG. 2004.** Roots, cycles and leaves. Expression of the phosphoenolpyruvate carboxylase kinase gene family in soybean. *Plant Physiology*, **135**, 2078-2087.
- Sung D-Y, Kaplan F, Lee K-J, Guy CL. 2003.** Acquired tolerance to temperature extremes. *Trends in Plant Science*, **8**, 179-187.
- Szecowka M, Heise R, Tohge T, Nunes-Nesi A, Vosloh D, Huege J, Feil R, Lunn J, Nikoloski Z, Stitt M, Fernie AR, Arrivault S. 2013.** Metabolic fluxes in an illuminated *Arabidopsis* rosette. *The Plant Cell*, **25**, 694.



- Tcherkez G, Gauthier P, Buckley TN, Busch FA, Barbour MM, Bruhn D, Heskell MA, Gong XY, Crous KY, Griffin K, et al. 2017.** Leaf day respiration: low CO<sub>2</sub> flux but high significance for metabolism and carbon balance. *New Phytologist*, **216**, 986-1001.
- Tcherkez G, Mahé A, Gauthier P, Mauve C, Gout E, Bligny R, Cornic G, Hodges M. 2009.** In folio respiratory fluxomics revealed by <sup>13</sup>C isotopic labeling and h/d isotope effects highlight the noncyclic nature of the tricarboxylic acid “cycle” in illuminated leaves. *Plant Physiology*, **151**, 620-630.
- Tovar-Mendez A, Miernyk JA, Randall DD. 2003.** Regulation of pyruvate dehydrogenase complex activity in plant cells. *Eur J Biochem*, **270**, 1043-1049.
- Urbanczyk-Wochniak E, Baxter C, Kolbe A, Kopka J, Sweetlove LJ, Fernie AR. 2005.** Profiling of diurnal patterns of metabolite and transcript abundance in potato (*Solanum tuberosum*) leaves. *Planta*, **221**, 891-903.
- Vanderwel MC, Slot M, Lichstein JW, Reich PB, Kattge J, Atkin OK, Bloomfield KJ, Tjoelker MG, Kitajima K. 2015.** Global convergence in leaf respiration from estimates of thermal acclimation across time and space. *New Phytologist*, **207**, 1026-1037.
- Walker BJ, VanLoocke A, Bernacchi CJ, Ort DR. 2016.** The cost of photorespiration to food production now and in the future. *Annual Review of Plant Biology*, **67**, 107-129.
- Wassmann R, Jagadish SVK, Heuer S, Ismail A, Redona E, Serraj R, Singh RK, Howell G, Pathak H, Sumfleth K. 2009.** Chapter 2: Climate change affecting rice production: The physiological and agronomic basis for possible adaptation strategies. In: *Advances in Agronomy*. (ed Sparks DL) pp 59-122. Academic Press.
- Watanabe CK, Sato S, Yanagisawa S, Uesono Y, Terashima I, Noguchi K. 2014.** Effects of elevated CO<sub>2</sub> on levels of primary metabolites and transcripts of genes encoding respiratory enzymes and their diurnal patterns in *Arabidopsis thaliana*: Possible relationships with respiratory rates. *Plant and Cell Physiology*, **55**, 341-357.
- Welch JR, Vincent JR, Auffhammer M, Moya PF, Dobermann A, Dawe D. 2010.** Rice yields in tropical/subtropical Asia exhibit large but opposing sensitivities to minimum and maximum temperatures. *Proceedings of the National Academy of Sciences*, **107**, 14562-14567.
- Werner RA, Buchmann N, Siegwolf RTW, Kornell BE, Gessler A. 2011.** Metabolic fluxes, carbon isotope fractionation and respiration – lessons to be learned from plant biochemistry. *New Phytologist*, **191**, 10-15.

- Weston DJ, Gunter LE, Rogers A, Wullschleger SD. 2008.** Connecting genes, coexpression modules, and molecular signatures to environmental stress phenotypes in plants. *BMC Systems Biology*, **2**, 16.
- Whitehead D, Griffin KL, Turnbull MH, Tissue DT, Engel VC, Brown KJ, Schuster WSF, Walcroft AS. 2004.** Response of total night-time respiration to differences in total daily photosynthesis for leaves in a *Quercus rubra* L. canopy: implications for modelling canopy CO<sub>2</sub> exchange. *Glob Change Biology*, **10**.
- Wickham H. 2017.** Tidyverse: Easily install and load the 'Tidyverse'. R package version 1.2.1.
- Zaragoza-Castells J, Sánchez-Gómez D, Valladares F, Hurry V, Atkin OK. 2007.** Does growth irradiance affect temperature dependence and thermal acclimation of leaf respiration? Insights from a Mediterranean tree with long-lived leaves. *Plant, Cell & Environment*, **30**, 820-833.
- Zell MB, Fahnenstich H, Maier A, Saigo M, Voznesenskaya EV, Edwards GE, Andreo C, Schleifenbaum F, Zell C, Drincovich MF, Maurino VG. 2010.** Analysis of Arabidopsis with highly reduced levels of malate and fumarate sheds light on the role of these organic acids as storage carbon molecules. *Plant Physiology*, **152**, 1251.
- Zhang B, Horvath S. 2005.** A general framework for weighted gene co-expression network analysis. *Statistical applications in genetics and molecular biology*, **4**.

## Supporting Information

**Dataset S1.** Metabolite profiles quantified by Gas Chromatography-Mass Spectrometry (GC-MS) for leaf samples collected for each growth treatment and over a diel cycle.

**Fig. S1** Photograph of rice plants in their hydroponic containers just prior to sampling.

Table S1. Two-way ANOVA of growth temperature and time effects on dark respiration measured at the prevailing day-time growth temperature and at a common temperature of 30°C.

Table S2. Two-way ANOVA of temperature and diurnal effects on the content of glucose, fructose, sucrose, total soluble sugar, starch and nitrogen in rice leaves.

Table S3. Results of linear regression on dark respiration rates ( $R_{\text{dark}}$ ) measured at 30°C and key metabolites identified in weighted correlation network analysis as presented in Figure 4.

### Figure legends

**Figure 1.** Dark respiration ( $R_{\text{dark}}$ ) of rice leaves on a dry mass basis measured over the diurnal cycle at; (a) the respective day-time growth temperature of each treatment; and (b), a common temperature of 30°C. Measurements were conducted on rice leaves grown at 25/20°C (blue squares), 30/25°C (control; green circles) and 40/35°C (red triangles). Area shaded in grey represents the night-time. The  $P$ -values from a two-way ANOVA for the temperature (Temp.), time of sampling (Time) and the interaction between the two variables (Temp.  $\times$  Time) are presented on the graphs. The  $F$ -values of the two-way ANOVA are presented in Supporting Information Table S1. Data represents means of four biological replicates  $\pm$  SE.

**Figure 2.** Rice leaf content of glucose (a), fructose (b), sucrose (c), total soluble sugar (d), starch (e), and nitrogen (f). Leaves were grown at different temperatures of 25/20°C (blue squares), 30/25°C (control; green circles) and 40/35°C (red triangles). Area shaded in grey represents the night-time. The  $P$ -values from a two-way ANOVA for the temperature (Temp.), time of sampling (Time) and the interaction between the two variables (Temp.  $\times$  Time) are presented on the graphs. The  $F$ -values of the two-way ANOVA are presented in Supporting Information Table S2. Data represents mean of three biological replicates  $\pm$  SE.

**Figure 3.** Principal component analyses (PCA) portraying relationships among metabolites in relation to the diurnal cycle and growth temperature. (a) Each thin arrow represents a metabolite, and for each arrow, the direction and length from the centre represent a correlation with principle components (x- and y-axis) and the quality of representation (loading) on the principle component plane. For example, Xylitol (labelled as 29) is well represented by both first and second principle components and is more correlated with the first than the second, while putrescine (labelled as 30) is not well represented by the first two components but is more correlated with the second than the first. Different colours of metabolites and their associated arrows denote different metabolite classes; amino acids in red, carbohydrates in green, organic acids in blue, and others in purple. (b) Metabolite responses in the first and

second dms of rice leaves to sampling time (day/night), with day indicated in red and night in blue. (c) Metabolite responses in the first and second dms of rice leaves to growth temperature with 25/20°C in blue, 30/25°C in green and 40/35°C in red.


**Figure 4.** Effect of the diurnal cycle on the relative concentrations of metabolites in rice leaves associated with carbohydrates, glycolysis, amino acids and the tricarboxylic acid (TCA) cycle. Rice leaves were sampled at five different time points throughout light/dark cycle: 13:00 (red), 17:30 (yellow), 21:30 (green), 04:30 (blue) and 08:30 (purple) hours. Data represents mean of four biological replicates. One-way ANOVA and post-hoc test were conducted and different letters show significant differences according to Tukey test ( $P < 0.05$ ).

**Figure 5.** Effect of growth temperature treatments on the relative abundance levels of metabolites in rice leaves associated with carbohydrate metabolism, glycolysis, amino acid metabolism and tricarboxylic acid (TCA) cycle. Measurements were conducted on rice leaves grown in 25/20°C (blue), 30/25°C (green) and 40/35°C (red). Data represents mean of four biological replicates. One-way ANOVA and post hoc test were conducted and different letters show significant differences according to Tukey test ( $P < 0.05$ ).


**Figure 6.** Metabolites that significantly correlated with  $R_{\text{dark}}$  at 30°C identified through weighted correlation network analysis. For aconitate, xylitol and erythritol, correlations were mostly attributable to growth temperature. For serine, putrescine and fructose-6-phosphate, correlations were attributable to both growth temperature and the diel cycle. Succinate correlation was attributable to the day/night cycle. Regression analysis  $P$  and  $r^2$  values for each metabolite are presented in Supporting Information Table S3.

**Table 1.** Two-way ANOVA of temperature and diurnal effects on the metabolite level of rice leaves grown at different temperatures. Asterisks (\*) denote statistically significant difference ( $P < 0.05$ ) among the levels of temperature treatments and time points.


Metabolite	<i>P</i> -value		
	Temperature	Diurnal	Temperature x Diurnal
Alanine	0.1130	$5.25 \times 10^{-7}$ *	0.2665
Valine	0.2452	0.0242*	0.4354
Isoleucine	0.0281*	0.2110	0.3534
Proline	0.0116*	0.7936	0.2217
Succinate	0.0108*	$7.28 \times 10^{-9}$ *	0.6848
Fumarate	0.0363*	0.0007*	0.9844
Threonine	0.9219	0.0747	0.1243
Malate	0.1522	$2.22 \times 10^{-13}$ *	0.2691
Aspartic acid	0.0829	$1.07 \times 10^{-6}$ *	0.2295
4-Aminobutyric acid	0.0028*	0.9579	0.5610
$\alpha$ -ketoglutarate	$3.32 \times 10^{-5}$ *	$< 2.2 \times 10^{-16}$ *	0.1675
Phenylalanine	$3.54 \times 10^{-6}$ *	0.0003*	0.0741
Asparagine	0.9157	0.2746	0.2260
Putrescine	$2.37 \times 10^{-5}$ *	0.1720	0.4763
Aconitate	$3.75 \times 10^{-9}$ *	0.0016*	0.6774
Glutamine	$1.04 \times 10^{-5}$ *	0.0279*	0.3447
Shikimic acid	0.5227	0.1504	0.4783
Citrate	0.0919	$8.19 \times 10^{-5}$ *	0.7641
Fructose	0.1620	0.1418	0.8123
Glucose	0.4284	0.0189*	0.9153
Lysine	$1.36 \times 10^{-6}$ *	$2.04 \times 10^{-5}$ *	0.4090
Tyrosine	$2.01 \times 10^{-8}$ *	$5.83 \times 10^{-12}$ *	0.0092*
Tryptophan	0.2963	$1.35 \times 10^{-5}$ *	0.0166*
Fructose-6-phosphate	$4.74 \times 10^{-14}$	$1.59 \times 10^{-7}$ *	0.5368
Glucose-6-phosphate	$5.62 \times 10^{-12}$	$5.18 \times 10^{-6}$ *	0.9891
Sucrose	$2.66 \times 10^{-5}$ *	0.0287*	0.7049



**Figure 1.** Dark respiration ( $R_{\text{dark}}$ ) of rice leaves on a dry mass basis measured over the diurnal cycle at; (a) the respective day-time growth temperature of each treatment; and (b), a common temperature of 30°C. Measurements were conducted on rice leaves grown at 25/20°C (blue squares), 30/25°C (control; green circles) and 40/35°C (red triangles). Area shaded in grey represents the night-time. The  $P$ -values from a two-way ANOVA for the temperature (Temp.), time of sampling (Time) and the interaction between the two variables (Temp. x Time) are presented on the graphs. The  $F$ -values of the two-way ANOVA are presented in Supporting Information Table S1. Data represents means of four biological replicates  $\pm$  SE.




**Figure 2.** Rice leaf content of glucose (a), fructose (b), sucrose (c), total soluble sugar (d), starch (e), and nitrogen (f). Leaves were grown at different temperatures of 25/20°C (blue squares), 30/25°C (control; green circles) and 40/35°C (red triangles). Area shaded in grey represents the night-time. The *P*-values from a two-way ANOVA for the temperature (Temp.), time of sampling (Time) and the interaction between the two variables (Temp.  $\times$  Time) are presented on the graphs. The *F*-values of the two-way ANOVA are presented in Supporting Information Table S2. Data represents mean of three biological replicates  $\pm$  SE.




**Figure 3.** Principal component analyses (PCA) portraying relationships among metabolites in relation to the diurnal cycle and growth temperature. (a) Each thin arrow represents a metabolite, and for each arrow, the direction and length from the centre represent a correlation with principle components (x- and y-axis) and the quality of representation (loading) on the principle component plane. For example, Xylitol (labelled as 29) is well represented by both first and second principle components and is more correlated with the first than the second, while putrescine (labelled as 30) is not well represented by the first two components but is more correlated with the second than the first. Different colours of metabolites and their associated arrows denote different metabolite classes; amino acids in red, carbohydrates in green, organic acids in blue, and others in purple. (b) Metabolite responses in the first and second dims of rice leaves to sampling time (day/night), with day indicated in red and night in blue. (c) Metabolite responses in the first and second dims of rice leaves to growth temperature with 25/20°C in blue, 30/25°C in green and 40/35°C in red.






**Figure 4.** Effect of the diurnal cycle on the relative concentrations of metabolites in rice leaves associated with carbohydrates, glycolysis, amino acids and the tricarboxylic acid (TCA) cycle. Rice leaves were sampled at five different time points throughout light/dark cycle: 13:00 (red), 17:30 (yellow), 21:30 (green), 04:30 (blue) and 08:30 (purple) hours. Data represents mean of four biological replicates. One-way ANOVA and post-hoc test were conducted and different letters show significant differences according to Tukey test ( $P < 0.05$ ).



**Figure 5.** Effect of growth temperature treatments on the relative abundance levels of metabolites in rice leaves associated with carbohydrate metabolism, glycolysis, amino acid metabolism and tricarboxylic acid (TCA) cycle. Measurements were conducted on rice leaves grown in 25/20°C (blue), 30/25°C (green) and 40/35°C (red). Data represents mean of four biological replicates. One-way ANOVA and post hoc test were conducted and different letters show significant differences according to Tukey test ( $P < 0.05$ ).



**Figure 6.** Metabolites that significantly correlated with  $R_{\text{dark}}$  at 30°C identified through weighted correlation network analysis. For aconitate, xylitol and erythritol, correlations were mostly attributable to growth temperature. For serine, putrescine and fructose-6-phosphate, correlations were attributable to both growth temperature and the diel cycle. Succinate correlation was attributable to the day/night cycle. Regression analysis  $P$  and  $r^2$  values for each metabolite are presented in Supporting Information Table S3.